



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Reply to: Oncolytic Viral Therapy for Malignant Pleural Mesothelioma

Jean, Didier ; Delaunay, Tiphaine ; Meiller, Clément ; Boisgerault, Nicolas ; Grard, Marion ; Caruso, Stefano ; Blanquart, Christophe ; Felley-Bosco, Emanuela ; Bennouna, Jaafar ; Tangy, Frédéric ; Grégoire, Marc ; Fonteneau, Jean-François

DOI: <https://doi.org/10.1016/j.jtho.2020.04.011>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-193570>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Jean, Didier; Delaunay, Tiphaine; Meiller, Clément; Boisgerault, Nicolas; Grard, Marion; Caruso, Stefano; Blanquart, Christophe; Felley-Bosco, Emanuela; Bennouna, Jaafar; Tangy, Frédéric; Grégoire, Marc; Fonteneau, Jean-François (2020). Reply to: Oncolytic Viral Therapy for Malignant Pleural Mesothelioma. *Journal of Thoracic Oncology*, 15(7):e113-e116.

DOI: <https://doi.org/10.1016/j.jtho.2020.04.011>

Journal of Thoracic Oncology

Reply to: Oncolytic Viral Therapy for Malignant Pleural Mesothelioma

--Manuscript Draft--

Manuscript Number:	
Full Title:	Reply to: Oncolytic Viral Therapy for Malignant Pleural Mesothelioma
Article Type:	Response to Letter to the Editor
Keywords:	
Corresponding Author:	Jean-François FONTENEAU, Ph.D INSERM Nantes, FRANCE
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	INSERM
Corresponding Author's Secondary Institution:	
First Author:	Jean-François FONTENEAU, Ph.D
First Author Secondary Information:	
Order of Authors:	Jean-François FONTENEAU, Ph.D
	Didier Jean, PhD
	Tiphaine Delaunay, PhD
	Clément Meiller, MS
	Nicolas Boisgerault, PhD
	Marion Grard, MS
	Stefano Caruso, PhD
	Christophe Blanquart, PhD
	Emanuela Felley-Bosco, PhD
	Jaafar Bennouna, MD
	Frédéric Tangy, PhD
	Marc Grégoire, PhD
Order of Authors Secondary Information:	
Manuscript Region of Origin:	FRANCE

Reply to : Oncolytic Viral Therapy for Malignant Pleural Mesothelioma

Didier Jean¹, Tiphaine Delaunay^{2,3}, Clément Meiller¹, Nicolas Boisgerault^{2,3}, Marion Grard^{2,3}, Stefano Caruso¹, Christophe Blanquart^{2,3}, Emanuela Felley-Bosco⁴, Jaafar Bennouna^{2,3,5}, Frédéric Tangy⁶, Marc Grégoire^{2,3}, and Jean-François Fonteneau^{2,3}

¹ Centre de Recherche des Cordeliers, Inserm, Sorbonne Université, Université de Paris, Functional Genomics of Solid Tumors, F-75006 Paris, France.

² CRCINA, INSERM, CNRS, Université d'Angers, Université de Nantes, Nantes, France

³ Labex IGO, Immunology Graft Oncology, Nantes, France

⁴ Laboratory of Molecular Oncology Lungen- und Thoraxonkologie Zentrum Zurich University Hospital

⁵ CHU de Nantes, oncologie thoracique et digestive, Université de Nantes, 44000, France

⁶ CNRS 3569, Institut Pasteur, Paris, 75015, France

Running title: *BAP1* status and the type I interferon pathway in mesothelioma

Key words: Oncolytic immunotherapy, measles virus, type I interferon, gene homozygous deletion, mesothelioma.

Financial supports : This work was supported by “La Ligue Régionale Grand Ouest contre le Cancer” (CSIRGO: CD16, CD22, CD44, CD49, CD72, CD79 and CD85), Ile de France committee of “Ligue contre le Cancer”, “La Ligue Nationale contre le Cancer”, “L’association ARSMESO44”, “La Fondation du Souffle et le Fonds de Dotation Recherche en Santé Respiratoire”, “La fondation ARC”, “La Fondation pour la Recherche Médicale (FRM)”, L’Agence Nationale pour la Recherche (ANR-16-CE18-0016), “bourse Roche de Recherche 2015 en Oncologie Thoracique”, INSERM, “LabEX IGO program supported by the National Research Agency via the investment of the future program ANR-11-LABX-0016-01”, Hadassah France and Chancellerie des Universités de Paris (Legs POIX). TD was supported by a grant from Ligue contre le Cancer.

Corresponding authors: Jean-François Fonteneau, INSERM U1232, CRCINA, Institut de Recherche en Santé de l'Université de Nantes, 8 quai Moncousu, BP70721, 44007 Nantes cedex 1, France. Email : jean-francois.fonteneau@inserm.fr
Didier Jean, Centre de Recherche des Cordeliers / Inserm UMRS 1138 (ex Inserm UMR 1162), Equipe « Génomique Fonctionnelle des Tumeurs Solides » - FunGeST 15 rue de l'Ecole de Médecine, 75006 PARIS, France. Email : didier.jean@inserm.fr

787 words, 1 figure, 8 references

1 Among the most frequent genomic alterations in malignant pleural
2 mesothelioma (MPM) are the homozygous deletions (HD) of the *CDKN2A* tumor
3 suppressor gene. These alterations are found in 49% of MPM patients according to a
4 recent analysis of TCGA, The Cancer Genome Atlas ¹. In our recent study published
5 in the Journal of Thoracic Oncology performed on 78 short-term-cultured MPM cell
6 lines, we found *CDKN2A* HD in 73% of patients ². We also showed that the *CDKN2A*
7 HD are accompanied in 30% of them by the HD of all genes encoding type I interferons
8 (IFN I) that are essential for cellular antiviral defense. Finally, we identified two MPM
9 cell lines with a constitutively activated IFN I response characterized by the expression
10 of numerous IFN-stimulated genes (ISG) that resist to the oncolytic activity of the
11 vaccine Schwarz strain of Measles virus (MV).
12
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Other frequent genomic alterations in MPM are the mutations and/or the HD of
27 the *BAP1* gene ¹. In their letter to the editor, Yang H *et al* focus on The Cancer Genome
28 Atlas (TCGA) data and show that *BAP1* mutation and *BAP1* HD are found in 22% and
29 14% of patients respectively. They performed a functional enrichment analysis using
30 Gene Ontology and Reactome databases to identify the signal pathways that correlate
31 to *BAP1* expression. They show a strong negative correlation between *BAP1*
32 expression and a constitutively activated IFN I response. They conclude that *BAP1*
33 loss-of-function may be used as a marker of tumors with constitutively activated IFN I
34 response and thus resistance to oncolytic virotherapy.
35
36
37
38
39
40
41
42
43
44
45
46
47

48 Hmeljak J *et al* previously performed an analysis of TCGA on 74 MPM patients
49 with a detailed focus on *BAP1* alteration and its consequences ¹. They report an IFN I
50 signature in MPM tumors with an inactivated *BAP1* gene (Supplementary Figure 3H
51 and Supplementary Figure 7D). Thus, the expression of *BAP1*, as well as its mutational
52 status suggest a negative regulation of the IFN I pathway in MPM by BAP1 protein.
53
54
55
56
57
58
59
60
61
62
63
64
65

Considering those results, we first analyzed *BAP1* gene expression in 19 MPM cells lines for which we previously determined their MV-sensitivity ³. We observed a significant enrichment of low *BAP1*-expressing MPM cell lines that are MV-resistant (p=0.02) (**Figure 1A**). We then screened *BAP1* gene alterations in these cell lines. Eight MPM cell lines harbor *BAP1* alterations, 3 with HD and 5 with mutations (**Figure 1B**). These data show that all MPM cell lines with *IFNB1* deletion are sensitive to MV independently of the *BAP1* mutation status. Interestingly, a slight enrichment of *BAP1* genetic alterations was observed in MV-resistant compared to MV-sensitive MPM cell lines (p=0.07), especially considering MPM cell lines without *IFNB1* deletion (p=0.06) (**Figure 1C**).

We thus analyzed the IFN I signature of 36 short-term-cultured MPM cell lines for which *IFNB1* and *BAP1* genetic alteration status have been determined (61% with *BAP1*, 28% with *IFNB1*, 14% with both genes alteration). We performed a pathway enrichment analysis using transcriptomic data previously generated from these cell lines ⁴ and the online pathway analytic tool of Reactome Knowledgebase (<https://reactome.org/>). We did not find any significant enrichment of IFN I pathways on genes whose expression inversely correlates with *BAP1* gene expression (Spearman's $r < -0.4$, FDR-adjusted $p < 0.01$), neither on differentially expressed genes between MPM cell lines with or without *BAP1* genetic alteration (data not shown). This absence of link between *BAP1* expression or alteration and IFN I pathways was also observed in our previous study ². In absence of virus, MV-resistant Meso61 cell line with a bi-allelic deletion and low expression of *BAP1* harbors an inactivated IFN I response, while MV-resistant Meso45 with a high expression of wild-type *BAP1* harbors a strong constitutively activated IFN I.

1 Altogether, our results suggest a correlation between *BAP1* gene expression or
2 alteration status and the resistance to MV, but we cannot conclude on a link with the
3
4 IFN I response. This discrepancy of the results between tumor samples and cell lines
5
6 may be explained by the microenvironment. It was reported that *BAP1*-altered MPM
7
8 tumors show a different pattern of infiltration by immune cells than wild-type *BAP1*
9
10 tumors ⁵ and these infiltrations may play a role in the observed IFN I signature.
11
12 Furthermore, this IFN I signature may be responsible for the presence of infiltrated
13
14 immune cells as it was reported for other cancers ⁶.
15
16
17
18

19 In conclusion, the negative correlation of *BAP1* expression and the IFN I
20
21 signature in MPM observed by Yang H et al could have a great interest in the context
22
23 of absence of *INFB1* deletion to identify patient with a lower chance to respond to
24
25 oncolytic virotherapy using IFN I-sensitive oncolytic viruses. It would thus be interesting
26
27 to determine the mechanism of the IFN I response modulation by the BAP1 protein.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Reference:


1. Hmeljak J, Sanchez-Vega F, Hoadley KA, et al. Integrative Molecular Characterization of Malignant Pleural Mesothelioma. *Cancer Discov* 2018;8:1548-1565.
2. Delaunay T, Achard C, Boisgerault N, et al. Frequent Homozygous Deletions of Type I Interferon Genes in Pleural Mesothelioma Confer Sensitivity to Oncolytic Measles Virus. *J Thorac Oncol* 2020.
3. Achard C, Boisgerault N, Delaunay T, et al. Sensitivity of pleural mesothelioma to oncolytic measles virus depends on defects of the type I interferon response. *Oncotarget* 2015;Dec 29 6:44892-44904.
4. de Reynies A, Jaurand MC, Renier A, et al. Molecular classification of malignant pleural mesothelioma: identification of a poor prognosis subgroup linked to the epithelial-to-mesenchymal transition. *Clin Cancer Res* 2014;20:1323-1334.
5. Ladanyi M, Sanchez Vega F, Zauderer M. Loss of BAP1 as a candidate predictive biomarker for immunotherapy of mesothelioma. *Genome Med* 2019;11:18.
6. Canadas I, Thummalapalli R, Kim JW, et al. Tumor innate immunity primed by specific interferon-stimulated endogenous retroviruses. *Nat Med* 2018;24:1143-1150.
7. Blum Y, Meiller C, Quetel L, et al. Dissecting heterogeneity in malignant pleural mesothelioma through histo-molecular gradients for clinical applications. *Nat Commun* 2019;10:1333.
8. Quetel L, Meiller C, Assie JB, et al. Genetic alterations of malignant pleural mesothelioma: association with tumor heterogeneity and overall survival. *Mol Oncol* 2020.

Figure legends

Figure 1: Comparison of *BAP1* gene expression and genetic alteration status to Measles virus sensitivity in malignant pleural mesothelioma cell lines. *BAP1* gene expression and mutation status were analyzed in 19 malignant pleural mesothelioma (MPM) cell lines characterized for Measles virus (MV) sensitivity in our previous study ³. **A.** *BAP1* gene expression was determined by RT-qPCR as previously described ⁷. MV resistant cell lines showed a significantly lower *BAP1* expression than MV sensitive cell lines (Mann Whitney test, p-value=0.02). **B.** *BAP1* mutation status was defined using a targeted next generation sequencing ⁸. Biallelic deletions were detected based on sequencing coverage. Sequencing data analysis showed mutations in 8 cell lines: (i) 3 cell lines with bi-allelic deletions of *BAP1* exons, (ii) 2 with nonsense mutations, (iii) 2 with splice site mutations and (iv) 1 with missense mutation predicted to be functionally damaging by polyphen HDIV and SIFT. **C.** MV sensitivity and *INFB1* and *BAP1* mutations profile in MPM cell lines were resumed in a heat map. *BAP1* mutation and deletion are enriched in MV resistant cell lines compared to MV sensitive cell lines (Fisher's exact test, p=0.07 in all MPM cell lines and p=0.06 in MPM cell lines without *INFB1* deletion). Statistical tests were performed using GraphPad Prism version 6.07 software.

Figure 1

Figure 1



Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.